arXiv:math/0302276v5 [math.MG] 2 Oct 2006

A discrete form of the Beckman-Quarles theorem for mappings from $\mathbb{R}^2$ ($\mathbb{C}^2$) to $\mathbb{F}^2$, where $\mathbb{F}$ is a subfield of a commutative field extending $\mathbb{R}$ ($\mathbb{C}$)

Apoloniusz Tyszka

Abstract. Let $\mathbb{F}$ be a subfield of a commutative field extending $\mathbb{R}$. Let $\varphi_n : \mathbb{F}^n \times \mathbb{F}^n \to \mathbb{F}$, $\varphi_n((x_1, ..., x_n), (y_1, ..., y_n)) = (x_1 - y_1)^2 + ... + (x_n - y_n)^2$. We say that $f : \mathbb{R}^n \to \mathbb{F}^n$ preserves distance $d \geq 0$ if $|x - y| = d$ implies $\varphi_n(f(x), f(y)) = d^2$ for each $x, y \in \mathbb{R}^n$. Let $A_n(\mathbb{F})$ denote the set of all positive numbers d such that any map $f : \mathbb{R}^n \to \mathbb{F}^n$ that preserves unit distance preserves also distance d. Let $D_n(\mathbb{F})$ denote the set of all positive numbers d with the property: if $x, y \in \mathbb{R}^n$ and $|x - y| = d$ then there exists a finite set S_{xy} with $\{x, y\} \subseteq S_{xy} \subseteq \mathbb{R}^n$ such that any map $f : S_{xy} \to \mathbb{F}^n$ that preserves unit distance satisfies $|x - y|^2 = \varphi_n(f(y), f(y))$. Obviously, $\{1\} \subseteq D_n(\mathbb{F}) \subseteq A_n(\mathbb{F})$. We prove: $A_n(\mathbb{C}) \subseteq \left\{d > 0 : d^2 \in \mathbb{Q}\right\} \subseteq D_2(\mathbb{F})$. Let $\mathbb{K}$ be a subfield of a commutative field Γ extending $\mathbb{C}$. Let $\psi_2 : \Gamma^2 \times \Gamma^2 \to \Gamma$, $\psi_2((x_1, x_2), (y_1, y_2)) = (x_1 - y_1)^2 + (x_2 - y_2)^2$. We say that $f : \mathbb{C}^2 \to \mathbb{K}^2$ preserves unit distance if $\psi_2(X, Y) = 1$ implies $\psi_2(f(X), f(Y)) = 1$ for each $X, Y \in \mathbb{C}^2$. We prove: if $X, Y \in \mathbb{C}^2$, $\psi_2(X, Y) \in \mathbb{Q}$ and $X \neq Y$, then there exists a finite set S_{XY} with $\{X, Y\} \subseteq S_{XY} \subseteq \mathbb{C}^2$ such that any map $f : S_{XY} \to \mathbb{K}^2$ that preserves unit distance satisfies $\psi_2(X, Y) = \psi_2(f(X), f(Y))$ and $f(X) \neq f(Y)$.

2000 Mathematics Subject Classification: 51M05.

Key words: affine (semi-affine) isometry, affine (semi-affine) mapping with orthogonal linear part, Beckman-Quarles theorem, Cayley-Menger determinant, endomorphism (automorphism) of the field of complex numbers, unit-distance preserving mapping.

Let $\mathbb{F}$ be a subfield of commutative field extending $\mathbb{R}$. Let $\varphi_n : \mathbb{F}^n \times \mathbb{F}^n \to \mathbb{F}$, $\varphi_n((x_1, ..., x_n), (y_1, ..., y_n)) = (x_1 - y_1)^2 + ... + (x_n - y_n)^2$. We say that $f : \mathbb{R}^n \to \mathbb{F}^n$ preserves distance $d \geq 0$ if $|x - y| = d$ implies $\varphi_n(f(x), f(y)) = d^2$ for each $x, y \in \mathbb{R}^n$. We say that $f : \mathbb{R}^n \to \mathbb{F}^n$ preserves the distance between $x, y \in \mathbb{R}^n$ if $|x - y|^2 = \varphi_n(f(x), f(y))$. By a *complex isometry* of $\mathbb{C}^n$ (i.e. an affine isometry of $\mathbb{C}^n$) we understand any map $h : \mathbb{C}^n \to \mathbb{C}^n$ of the form

$$h(z_1, z_2, ..., z_n) = (z'_1, z'_2, ..., z'_n)$$

where

$$z'_j = a_{0j} + a_{1j}z_1 + a_{2j}z_2 + ... + a_{nj}z_n \qquad (j = 1, 2, ..., n),$$

the coefficients a_{ij} are complex and the matrix $||a_{ij}||$ $(i, j = 1, 2, ..., n)$ is orthogonal i.e. satisfies the condition

$$\sum_{j=1}^{n} a_{\mu j} a_{\nu j} = \delta_{\nu}^{\mu} \qquad (\mu, \nu = 1, 2, ..., n)$$

with Kronecker's delta. According to [6], $\varphi_n(x, y)$ is invariant under complex isometries i.e. for every complex isometry $h : \mathbb{C}^n \to \mathbb{C}^n$

$$\forall x, y \in \mathbb{C}^n \quad \varphi_n(h(x), h(y)) = \varphi_n(x, y) \tag{$\diamond$}$$

Conversely, if $h : \mathbb{C}^n \to \mathbb{C}^n$ satisfies $(\diamond)$ then h is a complex isometry; it follows from [4, proposition 1, page 21] by replacing $\mathbb{R}$ with $\mathbb{C}$ and $d(x, y)$ with $\varphi_n(x, y)$. Similarly, if $f : \mathbb{R}^n \to \mathbb{C}^n$ preserves all distances, then there exists a complex isometry $h : \mathbb{C}^n \to \mathbb{C}^n$ such that $f = h_{|\mathbb{R}^n}$.

By a field endomorphism of a field H we understand any ring endomorphism $g : H \to H$ with $g(1) \neq 0$. Bijective endomorphisms are called automorphisms, for more information on field endomorphisms and automorphisms of $\mathbb{C}$ the reader is referred to [8], [9] and [15]. If r is a rational number, then $g(r) = r$ for any field endomorphism $g : \mathbb{C} \to \mathbb{C}$. Proposition 1 shows that only rational numbers r have this property:

Proposition 1 ([12]). If $r \in \mathbb{C}$ and r is not a rational number, then there exists a field automorphism $g : \mathbb{C} \to \mathbb{C}$ such that $g(r) \neq r$.

Theorem 1 ([12]). If $x, y \in \mathbb{R}^n$ and $|x - y|^2$ is not a rational number, then there exists $f : \mathbb{R}^n \to \mathbb{C}^n$ that does not preserve the distance between x and y although f preserves all distances $\sqrt{r}$ with rational $r \geq 0$.

Let $A_n(\mathbb{F})$ denote the set of all positive numbers d such that any map $f : \mathbb{R}^n \to \mathbb{F}^n$ that preserves unit distance preserves also distance d. The classical Beckman-Quarles theorem states that each unit-distance preserving mapping from $\mathbb{R}^n$ to $\mathbb{R}^n$ $(n \geq 2)$ is an isometry, see [1]-[4] and [7]. It means that $A_n(\mathbb{R}) = (0, \infty)$ for each $n \geq 2$. By Theorem 1 $A_n(\mathbb{C}) \subseteq \{d > 0 : d^2 \in \mathbb{Q}\}$. Let $D_n(\mathbb{F})$ denote the set of all positive numbers d with the following property:

($\star$) if $x, y \in \mathbb{R}^n$ and $|x - y| = d$ then there exists a finite set S_{xy} with $\{x, y\} \subseteq S_{xy} \subseteq \mathbb{R}^n$ such that any map $f : S_{xy} \to \mathbb{F}^n$ that preserves unit distance preserves also the distance between x and y.

Obviously, $\{1\} \subseteq D_n(\mathbb{F}) \subseteq A_n(\mathbb{F})$. The author proved in [12] that $D_n(\mathbb{C})$ is a dense subset of $(0, \infty)$ for each $n \geq 2$. The proof remains valid for $D_n(\mathbb{F})$. It is also known ([10], [11]) that $D_n(\mathbb{R})$ is equal to the set of positive algebraic numbers for each $n \geq 2$. We shall prove that $\{d > 0 : d^2 \in \mathbb{Q}\} \subseteq D_2(\mathbb{F})$. We need the following technical Propositions 2-5.

Proposition 2 (cf. [5], [6]). The points

$$c_1 = (z_{1,1},\ \dots,\ z_{1,n}),\ \dots,\ c_{n+1} = (z_{n+1,1},\ \dots,\ z_{n+1,n}) \in \mathbb{F}^n$$

are affinely dependent if and only if their Cayley-Menger determinant

$$\Delta(c_1,\dots,c_{n+1}) := \det \begin{bmatrix} 0 & 1 & 1 & \dots & 1 \\ 1 & 0 & \varphi_n(c_1,c_2) & \dots & \varphi_n(c_1,c_{n+1}) \\ 1 & \varphi_n(c_2,c_1) & 0 & \dots & \varphi_n(c_2,c_{n+1}) \\ \dots & \dots & \dots & \dots & \dots \\ 1 & \varphi_n(c_{n+1},c_1) & \varphi_n(c_{n+1},c_2) & \dots & 0 \end{bmatrix}$$

equals 0.

Proof. It follows from the equality

$$\left(\det \begin{bmatrix} z_{1,1} & z_{1,2} & \dots & z_{1,n} & 1 \\ z_{2,1} & z_{2,2} & \dots & z_{2,n} & 1 \\ \dots & \dots & \dots & \dots & \dots \\ z_{n+1,1} & z_{n+1,2} & \dots & z_{n+1,n} & 1 \end{bmatrix} \right)^2 = \frac{(-1)^{n+1}}{2^n} \cdot \Delta(c_1,\dots,c_{n+1}).$$

Proposition 3 (cf. [5], [6]). For any points $c_1,\dots,c_{n+k} \in \mathbb{F}^n$ $(k = 2,3,4,\dots)$ their Cayley-Menger determinant equals 0 i.e. $\Delta(c_1,\dots,c_{n+k}) = 0$.

Proof. Assume that $c_1 = (z_{1,1},\ \dots,\ z_{1,n}),\ \dots,\ c_{n+k} = (z_{n+k,1},\ \dots,\ z_{n+k,n})$. The points

$$\widetilde{c}_1 = (z_{1,1},\ \dots,\ z_{1,n},\ 0,\dots,0),\ \dots,\ \widetilde{c}_{n+k} = (z_{n+k,1},\ \dots,\ z_{n+k,n},\ 0,\dots,0) \in \mathbb{F}^{n+k-1}$$

are affinely dependent. Since $\varphi_n(c_i,c_j) = \varphi_{n+k-1}(\widetilde{c}_i,\widetilde{c}_j)$ $(1 \leq i \leq j \leq n+k)$ the Cayley-Menger determinant of points $c_1,\dots,c_{n+k}$ is equal to the Cayley-Menger determinant of points $\widetilde{c}_1,\dots,\widetilde{c}_{n+k}$ which equals 0 according to Proposition 2.

Proposition 4a. If $d \in \mathbb{F}$, $d \neq 0$, $c_1,c_2,c_3 \in \mathbb{F}^2$ and $\varphi_2(c_1,c_2) = \varphi_2(c_1,c_3) = \varphi_2(c_2,c_3) = d^2$, then c_1, c_2, c_3 are affinely independent.

Proof. It follows from Proposition 2 because the Cayley-Menger determinant

$$\Delta(c_1,c_2,c_3) = \det \begin{bmatrix} 0 & 1 & 1 & 1 \\ 1 & 0 & d^2 & d^2 \\ 1 & d^2 & 0 & d^2 \\ 1 & d^2 & d^2 & 0 \end{bmatrix} = -3d^4 \neq 0.$$

Proposition 4b ([13]). If $a, b \in \mathbb{F}$, $a+b \neq 0$, $z, x, \widetilde{x} \in \mathbb{F}^2$ and $\varphi_2(z,x) = a^2$, $\varphi_2(x,\widetilde{x}) = b^2$, $\varphi_2(z,\widetilde{x}) = (a+b)^2$, then $\overrightarrow{zx} = \frac{a}{a+b}\overrightarrow{z\widetilde{x}}$.

Proposition 5 (cf. [6, Lemma, page 127] for $\mathbb{R}^n$). If $x, y, c_0, c_1, c_2 \in \mathbb{F}^2$, $\varphi_2(x,c_0) = \varphi_2(y,c_0)$, $\varphi_2(x,c_1) = \varphi_2(y,c_1)$, $\varphi_2(x,c_2) = \varphi_2(y,c_2)$ and c_0, c_1, c_2 are affinely independent, then $x = y$.

Proof. Computing we obtain that the vector $\overrightarrow{xy}$ is perpendicular to each of the linearly independent vectors $\overrightarrow{c_0c_1}$, $\overrightarrow{c_0c_2}$. Thus the vector $\overrightarrow{xy}$ is perpendicular to every linear combination of vectors $\overrightarrow{c_0c_1}$ and $\overrightarrow{c_0c_2}$. In particular, the vector $\overrightarrow{xy}$ is perpendicular to each of the vectors $[1,0]$ and $[0,1]$. Therefore $\overrightarrow{xy} = 0$ and the proof is completed.

Lemma 1. If $d \in D_2(\mathbb{F})$ then $\sqrt{3} \cdot d \in D_2(\mathbb{F})$.

Proof. Let $d \in D_2(\mathbb{F})$, $x, y \in \mathbb{R}^2$ and $|x-y| = \sqrt{3} \cdot d$. Using the notation of Figure 1 we show that

$$S_{xy} := S_{y\widetilde{y}} \cup \bigcup_{i=1}^{2} S_{xp_i} \cup \bigcup_{i=1}^{2} S_{yp_i} \cup S_{p_1p_2} \cup \bigcup_{i=1}^{2} S_{x\widetilde{p}_i} \cup \bigcup_{i=1}^{2} S_{\widetilde{y}\widetilde{p}_i} \cup S_{\widetilde{p_1}\widetilde{p_2}}$$

satisfies condition $(\star)$.



Figure 1

$|x-y| = |x-\widetilde{y}| = \sqrt{3} \cdot d$, $|y-\widetilde{y}| = |p_1-p_2| = |\widetilde{p_1}-\widetilde{p_2}| = d$

$|x-p_i| = |y-p_i| = |x-\widetilde{p_i}| = |\widetilde{y}-\widetilde{p_i}| = d$ $(i = 1,2)$

Assume that $f : S_{xy} \to \mathbb{F}^2$ preserves unit distance. Since

$$S_{xy} \supseteq S_{y\widetilde{y}} \cup \bigcup_{i=1}^{2} S_{xp_i} \cup \bigcup_{i=1}^{2} S_{yp_i} \cup S_{p_1p_2}$$

we conclude that f preserves the distances between y and $\widetilde{y}$, x and p_i $(i = 1,2)$, y and p_i $(i = 1,2)$, p_1 and p_2. Hence $\varphi_2(f(y), f(\widetilde{y})) = \varphi_2(f(x), f(p_i)) =$

$\varphi_2(f(y), f(p_i)) = \varphi_2(f(p_1), f(p_2)) = d^2$ $(i = 1, 2)$. By Proposition 3 the Cayley-Menger determinant $\Delta(f(x), f(p_1), f(p_2), f(y))$ equals 0 i.e.

$$\det \begin{bmatrix} 0 & 1 & 1 & 1 & 1 \\ 1 & 0 & \varphi_2(f(x), f(p_1)) & \varphi_2(f(x), f(p_2)) & \varphi_2(f(x), f(y)) \\ 1 & \varphi_2(f(p_1), f(x)) & 0 & \varphi_2(f(p_1), f(p_2)) & \varphi_2(f(p_1), f(y)) \\ 1 & \varphi_2(f(p_2), f(x)) & \varphi_2(f(p_2), f(p_1)) & 0 & \varphi_2(f(p_2), f(y)) \\ 1 & \varphi_2(f(y), f(x)) & \varphi_2(f(y), f(p_1)) & \varphi_2(f(y), f(p_2)) & 0 \end{bmatrix} = 0.$$

Therefore

$$\det \begin{bmatrix} 0 & 1 & 1 & 1 & 1 \\ 1 & 0 & d^2 & d^2 & t \\ 1 & d^2 & 0 & d^2 & d^2 \\ 1 & d^2 & d^2 & 0 & d^2 \\ 1 & t & d^2 & d^2 & 0 \end{bmatrix} = 0$$

where $t = \varphi_2(f(x), f(y))$. Computing this determinant we obtain

$$2d^2 t \cdot (3d^2 - t) = 0.$$

Therefore

$$t = \varphi_2(f(x), f(y)) = \varphi_2(f(y), f(x)) = 3d^2 = |x - y|^2$$

or

$$t = \varphi_2(f(x), f(y)) = \varphi_2(f(y), f(x)) = 0.$$

Analogously we may prove that

$$\varphi_2(f(x), f(\widetilde{y})) = \varphi_2(f(\widetilde{y}), f(x)) = 3d^2 = |x - \widetilde{y}|^2$$

or

$$\varphi_2(f(x), f(\widetilde{y})) = \varphi_2(f(\widetilde{y}), f(x)) = 0.$$

If $t = 0$ then the points $f(x)$ and $f(y)$ satisfy:

$$\varphi_2(f(x), f(x)) = 0 = \varphi_2(f(y), f(x)),$$
$$\varphi_2(f(x), f(p_1)) = d^2 = \varphi_2(f(y), f(p_1)),$$
$$\varphi_2(f(x), f(p_2)) = d^2 = \varphi_2(f(y), f(p_2)).$$

By Proposition 4a the points $f(x)$, $f(p_1)$, $f(p_2)$ are affinely independent. Therefore by Proposition 5 $f(x) = f(y)$ and consequently

$$d^2 = \varphi_2(f(y), f(\widetilde{y})) = \varphi_2(f(x), f(\widetilde{y})) \in \{3d^2,\ 0\}.$$

Since $d^2 \neq 3d^2$ and $d^2 \neq 0$ we conclude that the case $t = 0$ cannot occur. This completes the proof of Lemma 1.

Lemma 2. If $d \in D_2(\mathbb{F})$ then $2 \cdot d \in D_2(\mathbb{F})$.

Proof. Let $d \in D_2(\mathbb{F})$, $x, y \in \mathbb{R}^2$ and $|x - y| = 2 \cdot d$. Using the notation of Figure 2 we show that

$$S_{xy} := \bigcup \{S_{ab} : a, b \in \{x, y, p_1, p_2, p_3\}, |a - b| = d \vee |a - b| = \sqrt{3} \cdot d\}$$

(where S_{xp_3} and S_{yp_2} are known to exist by Lemma 1) satisfies condition $(\star)$.



Figure 2

$$|x-y| = 2 \cdot d$$

$$|p_1-p_2| = |p_1-p_3| = |p_2-p_3| = |x-p_1| = |x-p_2| = |y-p_1| = |y-p_3| = d$$

$$|x-p_3| = |y-p_2| = \sqrt{3} \cdot d$$

Assume that $f : S_{xy} \rightarrow \mathbb{F}^2$ preserves unit distance. Then f preserves all distances between p_i and p_j $(1 \leq i < j \leq 3)$, x and p_i $(1 \leq i \leq 3)$, y and p_i $(1 \leq i \leq 3)$. By Proposition 3 the Cayley-Menger determinant $\Delta(f(x),\ f(p_1),\ f(p_2),\ f(p_3),\ f(y))$ equals 0 i.e.

$$\det \begin{bmatrix} 0 & 1 & 1 & 1 & 1 & 1 \\ 1 & 0 & \varphi_2(f(x),f(p_1)) & \varphi_2(f(x),f(p_2)) & \varphi_2(f(x),f(p_3)) & \varphi_2(f(x),f(y)) \\ 1 & \varphi_2(f(p_1),f(x)) & 0 & \varphi_2(f(p_1),f(p_2)) & \varphi_2(f(p_1),f(p_3)) & \varphi_2(f(p_1),f(y)) \\ 1 & \varphi_2(f(p_2),f(x)) & \varphi_2(f(p_2),f(p_1)) & 0 & \varphi_2(f(p_2),f(p_3)) & \varphi_2(f(p_2),f(y)) \\ 1 & \varphi_2(f(p_3),f(x)) & \varphi_2(f(p_3),f(p_1)) & \varphi_2(f(p_3),f(p_2)) & 0 & \varphi_2(f(p_3),f(y)) \\ 1 & \varphi_2(f(y),f(x)) & \varphi_2(f(y),f(p_1)) & \varphi_2(f(y),f(p_2)) & \varphi_2(f(y),f(p_3)) & 0 \end{bmatrix} = 0.$$

Therefore

$$\det \begin{bmatrix} 0 & 1 & 1 & 1 & 1 & 1 \\ 1 & 0 & d^2 & d^2 & 3d^2 & t \\ 1 & d^2 & 0 & d^2 & d^2 & d^2 \\ 1 & d^2 & d^2 & 0 & d^2 & 3d^2 \\ 1 & 3d^2 & d^2 & d^2 & 0 & d^2 \\ 1 & t & d^2 & 3d^2 & d^2 & 0 \end{bmatrix} = 0$$

where $t = \varphi_2(f(x), f(y))$. Computing this determinant we obtain

$$3d^4 \cdot (t-4d^2)^2 = 0.$$

Therefore

$$t = \varphi_2(f(x), f(y)) = \varphi_2(f(y), f(x)) = 4d^2 = |x-y|^2.$$

Lemma 3. If $a, b \in D_2(\mathbb{F})$ and $a > b$, then $\sqrt{a^2-b^2} \in D_2(\mathbb{F})$.

Proof. Let $a, b \in D_2(\mathbb{F})$, $a > b$, $x, y \in \mathbb{R}^2$ and $|x-y| = \sqrt{a^2-b^2}$. Using the notation of Figure 3 we show that

$$S_{xy} := S_{xp_1} \cup S_{xp_2} \cup S_{yp_1} \cup S_{yp_2} \cup S_{p_1p_2}$$

(where $S_{p_1p_2}$ is known to exist by Lemma 2) satisfies condition $(\star)$.



Figure 3
$|x-y| = \sqrt{a^2-b^2}$
$|x-p_1| = |x-p_2| = b$, $|y-p_1| = |y-p_2| = a$, $|p_1-p_2| = 2b$

Assume that $f : S_{xy} \to \mathbb{F}^2$ preserves unit distance. Then f preserves the distances between x and p_i $(i = 1, 2)$, y and p_i $(i = 1, 2)$, p_1 and p_2. By Proposition 3 the Cayley-Menger determinant $\Delta(f(x),\ f(p_1),\ f(p_2),\ f(y))$ equals 0 i.e.

$$\det \begin{bmatrix} 0 & 1 & 1 & 1 & 1 \\ 1 & 0 & \varphi_2(f(x), f(p_1)) & \varphi_2(f(x), f(p_2)) & \varphi_2(f(x), f(y)) \\ 1 & \varphi_2(f(p_1), f(x)) & 0 & \varphi_2(f(p_1), f(p_2)) & \varphi_2(f(p_1), f(y)) \\ 1 & \varphi_2(f(p_2), f(x)) & \varphi_2(f(p_2), f(p_1)) & 0 & \varphi_2(f(p_2), f(y)) \\ 1 & \varphi_2(f(y), f(x)) & \varphi_2(f(y), f(p_1)) & \varphi_2(f(y), f(p_2)) & 0 \end{bmatrix} = 0.$$

Therefore

$$\det \begin{bmatrix} 0 & 1 & 1 & 1 & 1 \\ 1 & 0 & b^2 & b^2 & t \\ 1 & b^2 & 0 & 4b^2 & a^2 \\ 1 & b^2 & 4b^2 & 0 & a^2 \\ 1 & t & a^2 & a^2 & 0 \end{bmatrix} = 0$$

where $t = \varphi_2(f(x), f(y))$. Computing this determinant we obtain

$$-8b^2 \cdot (t + b^2 - a^2)^2 = 0.$$

Therefore

$$t = a^2 - b^2 = |x-y|^2.$$

Lemma 4. All distances $\sqrt{n}$ $(n = 1, 2, 3, ...)$ belong to $D_2(\mathbb{F})$.

Proof. If $n \in \{2, 3, 4, ...\}$ and $\sqrt{n} \in D_2(\mathbb{F})$, then $\sqrt{n-1} = \sqrt{(\sqrt{n})^2 - 1^2} \in D_2(\mathbb{F})$; it follows from Lemma 3 and $1 \in D_2(\mathbb{F})$. On the other hand, by Lemma 2 all numbers $\sqrt{2^{2k}} = 2^k$ $(k = 0, 1, 2, ..)$ belong to $D_2(\mathbb{F})$. These two facts imply that all distances $\sqrt{n}$ $(n = 1, 2, 3, ...)$ belong to $D_2(\mathbb{F})$.

From Lemma 4 we obtain:

Lemma 5. For each $n \in \{1,2,3,...\}$ we have: $n = \sqrt{n^2} \in D_2(\mathbb{F})$.

Lemma 6. If $d \in D_2(\mathbb{F})$ then all distances $\frac{d}{k}$ $(k = 2,3,4,...)$ belong to $D_2(\mathbb{F})$.

Proof. Let $d \in D_2(\mathbb{F})$, $k \in \{2,3,4,...\}$, $x,y \in \mathbb{R}^2$ and $|x-y| = \frac{d}{k}$. We choose an integer $e \geq d$. Using the notation of Figure 4 we show that

$$S_{xy} := S_{\widetilde{x}\widetilde{y}} \cup S_{zx} \cup S_{x\widetilde{x}} \cup S_{z\widetilde{x}} \cup S_{zy} \cup S_{y\widetilde{y}} \cup S_{z\widetilde{y}}$$

(where sets S_{zx}, $S_{x\widetilde{x}}$, $S_{z\widetilde{x}}$, S_{zy}, $S_{y\widetilde{y}}$, $S_{z\widetilde{y}}$ corresponding to integer distances are known to exist by Lemma 5) satisfies condition $(\star)$.



Figure 4
$|x-y| = \frac{d}{k}$

Assume that $f : S_{xy} \to \mathbb{F}^2$ preserves unit distance. Since $S_{xy} \supseteq S_{zx} \cup S_{x\widetilde{x}} \cup S_{z\widetilde{x}}$ we conclude that:

$$\varphi_2(f(z), f(x)) = |z-x|^2 = e^2,$$
$$\varphi_2(f(x), f(\widetilde{x})) = |x-\widetilde{x}|^2 = ((k-1)e)^2,$$
$$\varphi_2(f(z), f(\widetilde{x})) = |z-\widetilde{x}|^2 = (ke)^2.$$

By Proposition 4b:

$$\overrightarrow{f(z)f(x)} = \frac{1}{k}\overrightarrow{f(z)f(\widetilde{x})} \tag{1}$$

Analogously:

$$\overrightarrow{f(z)f(y)} = \frac{1}{k}\overrightarrow{f(z)f(\widetilde{y})} \tag{2}$$

By (1) and (2):

$$\overrightarrow{f(x)f(y)} = \frac{1}{k}\overrightarrow{f(\widetilde{x})f(\widetilde{y})} \tag{3}$$

Since $S_{xy} \supseteq S_{\widetilde{x}\widetilde{y}}$ we conclude that

$$\varphi_2(f(\widetilde{x}), f(\widetilde{y})) = |\widetilde{x}-\widetilde{y}|^2 = d^2 \tag{4}$$

By (3) and (4): $\varphi_2(f(x), f(y)) = \left(\frac{d}{k}\right)^2$ and the proof is completed.

Theorem 2. If $x,y \in \mathbb{R}^2$ and $|x-y|^2$ is a rational number, then there exists a finite set S_{xy} with $\{x,y\} \subseteq S_{xy} \subseteq \mathbb{R}^2$ such that any map $f : S_{xy} \to \mathbb{F}^2$ that preserves unit distance preserves also the distance between x and y; in other words $\{d > 0 : d^2 \in \mathbb{Q}\} \subseteq D_2(\mathbb{F})$.

Proof. We need to prove that $\sqrt{\frac{p}{q}} \in D_2(\mathbb{F})$ for all positive integers p, q. Since $\sqrt{\frac{p}{q}} = \frac{\sqrt{pq}}{q}$ the assertion follows from Lemmas 4 and 6.

Since $D_2(\mathbb{C}) \subseteq A_2(\mathbb{C}) \subseteq \{d > 0 : d^2 \in \mathbb{Q}\}$ as a corollary we get:

$$D_2(\mathbb{C}) = A_2(\mathbb{C}) = \{d > 0 : d^2 \in \mathbb{Q}\} \tag{$\bullet$}$$

Let $\mathbb{K}$ be a subfield of a commutative field Γ extending $\mathbb{C}$. Let $\psi_2 : \Gamma^2 \times \Gamma^2 \to \Gamma$, $\psi_2((x_1, x_2), (y_1, y_2)) = (x_1 - y_1)^2 + (x_2 - y_2)^2$. We say that $f : \mathbb{C}^2 \to \mathbb{K}^2$ preserves unit distance if $\psi_2(X, Y) = 1$ implies $\psi_2(f(X), f(Y)) = 1$ for each $X, Y \in \mathbb{C}^2$. We say that $f : \mathbb{C}^2 \to \mathbb{K}^2$ preserves the distance between $X, Y \in \mathbb{C}^2$ if $\psi_2(X, Y) = \psi_2(f(X), f(Y))$.

Theorem 3. If $X, Y \in \mathbb{C}^2$, $\psi_2(X, Y) \in \mathbb{Q}$ and $X \neq Y$, then there exists a finite set S_{XY} with $\{X, Y\} \subseteq S_{XY} \subseteq \mathbb{C}^2$ such that any map $f : S_{XY} \to \mathbb{K}^2$ that preserves unit distance satisfies $\psi_2(X, Y) = \psi_2(f(X), f(Y))$ and $f(X) \neq f(Y)$.

Proof. If $\psi_2(X, Y) = \psi_2(f(X), f(Y))$, then $\psi_2(X, Y) \neq 0$ implies $f(X) \neq f(Y)$. The main part of the proof divides into three cases.

Case 1: $\psi_2(X, Y) > 0$.

There exists a complex isometry of $\mathbb{C}^2$ that sends $(0, 0)$ to X and $(\sqrt{\psi_2(X, Y)}, 0)$ to Y, so without loss of generality we may assume that $X = (0, 0)$ and $Y = (\sqrt{\psi_2(X, Y)}, 0)$. By Theorem 2 there exists a finite set S_{XY} with $\{X, Y\} \subseteq S_{XY} \subseteq \mathbb{R}^2 \subseteq \mathbb{C}^2$ such that each unit-distance preserving mapping $f : S_{XY} \to \mathbb{K}^2$ satisfies $\psi_2(X, Y) = \psi_2(f(X), f(Y))$.

Case 2: $\psi_2(X, Y) < 0$.

Without loss of generality we may assume that $X = (0, 0)$. Let $Y = (a, b)$, $A = (-2ib, 2ia)$, $B = (2ib, -2ia)$. Then $a^2 + b^2 = \psi_2(X, Y) \in \mathbb{Q} \cap (-\infty, 0)$. Thus $\psi_2(A, X) = \psi_2(B, X) = -4(a^2 + b^2) \in \mathbb{Q} \cap (0, \infty)$, $\psi_2(A, B) = -16(a^2 + b^2) \in \mathbb{Q} \cap (0, \infty)$, $\psi_2(A, Y) = \psi_2(B, Y) = -3(a^2 + b^2) \in \mathbb{Q} \cap (0, \infty)$. We show that

$$S_{XY} := S_{AX} \cup S_{BX} \cup S_{AB} \cup S_{AY} \cup S_{BY}$$

satisfies the condition of the theorem; the finite sets S_{AX}, S_{BX}, S_{AB}, S_{AY}, S_{BY} are known to exist by the proof in case 1. Assume that $f : S_{XY} \to \mathbb{K}^2$ preserves unit distance. Then f preserves the distances between A and X, B and X, A and B, A and Y, B and Y. By Proposition 3 the Cayley-Menger determinant

$\Delta(f(X),f(Y),f(A),f(B))$ equals 0 i.e.

$$\det\begin{bmatrix} 0 & 1 & 1 & 1 & 1 \\ 1 & 0 & t & -4(a^2+b^2) & -4(a^2+b^2) \\ 1 & t & 0 & -3(a^2+b^2) & -3(a^2+b^2) \\ 1 & -4(a^2+b^2) & -3(a^2+b^2) & 0 & -16(a^2+b^2) \\ 1 & -4(a^2+b^2) & -3(a^2+b^2) & -16(a^2+b^2) & 0 \end{bmatrix} = 0$$

where $t = \psi_2(f(X),f(Y))$. Computing this determinant we obtain

$$32(a^2+b^2)(t-(a^2+b^2))^2 = 0.$$

Therefore $t = a^2+b^2 = \psi_2(X,Y)$.

Case 3: $\psi_2(X,Y) = 0$.

There exists a complex isometry of $\mathbb{C}^2$ that sends $(0,0)$ to X and $(1,i)$ to Y, so without loss of generality we may assume that $X = (0,0)$ and $Y = (1,i)$. Let $A = (-1,0)$, $B = (1,0)$. Then $\psi_2(A,X) = \psi_2(B,X) = 1$, $\psi_2(A,B) = 4$, $\psi_2(A,Y) = 3$, $\psi_2(B,Y) = -1$. We show that

$$S_{XY} := S_{AX} \cup S_{BX} \cup S_{AB} \cup S_{AY} \cup S_{BY}$$

satisfies the condition of the theorem; the finite sets S_{AX}, S_{BX}, S_{AB}, S_{AY}, S_{BY} are known to exist by the proofs in cases 1 and 2. Assume that $f : S_{XY} \to \mathbb{K}^2$ preserves unit distance. Then f preserves the distances between A and X, B and X, A and B, A and Y, B and Y. Since $1 = \psi_2(f(A),f(X)) \neq \psi_2(f(A),f(Y)) = 3$ we conclude that $f(X) \neq f(Y)$. By Proposition 3 the Cayley-Menger determinant $\Delta(f(X),f(Y),f(A),f(B))$ equals 0 i.e.

$$\det\begin{bmatrix} 0 & 1 & 1 & 1 & 1 \\ 1 & 0 & t & 1 & 1 \\ 1 & t & 0 & 3 & -1 \\ 1 & 1 & 3 & 0 & 4 \\ 1 & 1 & -1 & 4 & 0 \end{bmatrix} = 0$$

where $t = \psi_2(f(X),f(Y))$. Computing this determinant we obtain $-8t^2 = 0$. Therefore $t = 0 = \psi_2(X,Y)$.

The proof is finished.

As a corollary of Theorem 3 we get:

Theorem 4. If $X,Y \in \mathbb{C}^2$, $\psi_2(X,Y) \in \mathbb{Q}$ and $X \neq Y$, then any map $f : \mathbb{C}^2 \to \mathbb{K}^2$ that preserves unit distance satisfies $\psi_2(X,Y) = \psi_2(f(X),f(Y))$ and $f(X) \neq f(Y)$.

If $X,Y \in \mathbb{C}^2$ and $\psi_2(X,Y) \notin \mathbb{Q}$, then there exists $f : \mathbb{C}^2 \to \mathbb{C}^2$ that does not preserve the distance between X and Y although f satisfies $\psi_2(S,T) = \psi_2(f(S),f(T))$

for all $S, T \in \mathbb{C}^2$ with $\psi_2(S,T) \in \mathbb{Q}$; the proof is analogous to the proof of Theorem 1.

Using **(●)** the author proved ([13]):

(■) each unit-distance preserving mapping $f : \mathbb{R}^2 \to \mathbb{F}^2$ has a form $I \circ (\rho, \rho)$, where $\rho : \mathbb{R} \to \mathbb{F}$ is a field homomorphism and $I : \mathbb{F}^2 \to \mathbb{F}^2$ is an affine mapping with orthogonal linear part.

Using **(■)** and Theorem 4 the author proved ([14]):
each unit-distance preserving mapping $f : \mathbb{C}^2 \to \mathbb{C}^2$ has a form $I \circ (\gamma, \gamma)$, where $\gamma : \mathbb{C} \to \mathbb{C}$ is a field endomorphism and $I : \mathbb{C}^2 \to \mathbb{C}^2$ is an affine mapping with orthogonal linear part.

References

[1] F. S. Beckman and D. A. Quarles Jr., *On isometries of euclidean spaces*, Proc. Amer. Math. Soc. 4 (1953), 810–815.

[2] W. Benz, *An elementary proof of the theorem of Beckman and Quarles*, Elem. Math. 42 (1987), 4–9.

[3] W. Benz, *Geometrische Transformationen (unter besonderer Berücksichtigung der Lorentztransformationen)*, BI Wissenschaftsverlag, Mannheim, 1992.

[4] W. Benz, *Real geometries*, BI Wissenschaftsverlag, Mannheim, 1994.

[5] L. M. Blumenthal, *Theory and applications of distance geometry*, 2nd ed., Chelsea Publishing Co., New York, 1970.

[6] K. Borsuk, *Multidimensional analytic geometry*, Polish Scientific Publishers, Warsaw, 1969.

[7] U. Everling, *Solution of the isometry problem stated by K. Ciesielski*, Math. Intelligencer 10 (4) (1988), 47.

[8] H. Kestelman, *Automorphisms of the field of complex numbers*, Proc. Lond. Math. Soc., II. Ser. 53 (1951), 1–12.

[9] M. Kuczma, *An introduction to the theory of functional equations and inequalities: Cauchy's equation and Jensen's inequality*, Polish Scientific Publishers and Silesian University, Warsaw-Cracow-Katowice, 1985.

[10] A. Tyszka, *Discrete versions of the Beckman-Quarles theorem*, Aequationes Math. 59 (2000), 124–133.

[11] A. Tyszka, *Discrete versions of the Beckman-Quarles theorem from the definability results of R. M. Robinson*, Algebra, Geometry & their Applications, Seminar Proceedings 1 (2001), 88–90, Yerevan State University Press.

[12] A. Tyszka, *Beckman-Quarles type theorems for mappings from* $\mathbb{R}^n$ *to* $\mathbb{C}^n$, Aequationes Math. 67 (2004), 225–235.

[13] A. Tyszka, *The Beckman-Quarles theorem for mappings from* $\mathbb{R}^2$ *to* $\mathbb{F}^2$, *where* $\mathbb{F}$ *is a subfield of a commutative field extending* $\mathbb{R}$, Abh. Math. Sem. Univ. Hamburg 74 (2004), 77–87.

[14] A. Tyszka, *The Beckman-Quarles theorem for mappings from* $\mathbb{C}^2$ *to* $\mathbb{C}^2$, Acta Math. Acad. Paedagog. Nyházi. (N.S.) 21 (2005), no. 1, 63–69 (electronic).

[15] P. B. Yale, *Automorphisms of the complex numbers*, Math. Mag. 39 (1966), 135–141.

Apoloniusz Tyszka
Technical Faculty
Hugo Kołłątaj University
Balicka 116B, 30-149 Kraków, Poland
E-mail address: *rttyszka@cyf-kr.edu.pl*